

GAUGE INVARIANCE IN BOSON PRODUCTION

ROBIN G. STUART

*Randall Laboratory of Physics, University of Michigan, 500 E. University
Ann Arbor, Michigan 48109-1120, USA
E-mail: stuartr@umich.edu*

ABSTRACT

It is shown how to construct exactly gauge-invariant S -matrix elements for processes involving unstable gauge particles such as the W and Z^0 bosons. The results are applied to derive a physically meaningful expression for the cross-section $\sigma(e^+e^- \rightarrow Z^0 Z^0)$ and thereby provide a solution to the long-standing *problem of the unstable particle*. The problem of maintaining QED gauge-invariance in the process $e^+e^- \rightarrow \bar{\nu}_e e^- W^+ \rightarrow \bar{\nu}_e e^- u \bar{d}$ is examined.

1. Introduction

Pair production of W bosons has now been observed at LEP and both W 's and Z^0 's will be copiously produced at upcoming colliders. Yet both the W and Z^0 are unstable with quite sizable widths that cannot be treated using the standard methods of S -matrix theory because of the classic *problem of the unstable particle*¹. Roughly stated the problem is as follows: S -matrix theory deals with asymptotic in-states and out-states that propagate from and to infinity. Unstable particles cannot exist as asymptotic states because they decay a finite distance from the interaction region. Indeed it is known² that the S -matrix is unitary on the Hilbert space spanned by stable particle states and hence there is not even any room to accommodate unstable particles as external states. How can one use the S -matrix to calculate, say, the production cross-section for an unstable particle when it cannot exist as an asymptotic state?

In order to account for the finite width of the W and Z^0 one normally performs a Dyson summation of the one particle irreducible (1PI) self-energy diagrams that effectively replaces the tree-level propagator by a dressed propagator. However the 1PI self-energy is gauge-dependent starting at $\mathcal{O}(\alpha)$ and hence dressed propagator is gauge-dependent at all orders in α . If the dressed propagator is used in a finite-order calculation the resulting matrix element will also be gauge-dependent at some order. This gauge-dependence should be viewed as an indicator that the approximation scheme being used is inconsistent or does not represent a physical observable and is a latter-day manifestation of the problem of the unstable particle.

2. The Problem of the Unstable Particle

In this section we obtain a physical and therefore gauge-invariant property that

may be used to identify an unstable particle without requiring that it appear in the final state.

Recall that the coordinate space dressed propagator for a scalar particle has an integral representation

$$\Delta(x' - x) = \int \frac{d^4k}{(2\pi)^4} \frac{e^{-ik \cdot (x' - x)}}{k^2 - m^2 - \Pi(k^2) + i\epsilon} \quad (1)$$

The integrand has a pole at $k^2 = s_p$ where s_p is a solution of the equation $s - m^2 + \Pi(s) = 0$ and an analytic function, $F(s)$, via the relation $s - m^2 + \Pi(s) = (s - s_p)/F(s)$. The dressed propagator can then be written as

$$\Delta(x' - x) = \int \frac{d^4k}{(2\pi)^4} e^{-ik \cdot (x' - x)} \left[\frac{F(s_p)}{k^2 - s_p} + \frac{F(k^2) - F(s_p)}{k^2 - s_p} \right] \quad (2)$$

that separates resonant and non-resonant pieces. Performing the k_0 integration on the resonant part gives

$$\begin{aligned} \Delta(x' - x) = & -i \int \frac{d^3k}{(2\pi)^3 2k_0} e^{-ik \cdot (x' - x)} \theta(t' - t) F(s_p) \\ & + \int \frac{d^4k}{(2\pi)^4} \frac{F(k^2) - F(s_p)}{k^2 - s_p} - i \int \frac{d^3k}{(2\pi)^3 2k_0} e^{ik \cdot (x' - x)} \theta(t - t') F(s_p) \end{aligned}$$

where $k_0 = \sqrt{\vec{k}^2 + s_p}$. The non-resonant term contributes only for $t = t'$ and so represents a contact interaction. The resonant part splits into two terms that contribute when $t > t'$ or $t < t'$ and therefore connects points x and x' that are separated by a finite distance in space-time. When a similar analysis is applied to a physical matrix element one concludes that the resonant part describes a process in which there is a finite space-time separation between the initial-state vertex, V_i , and the final-state vertex, V_f . In other words, the resonant term describes the finite propagation of a physical unstable particle. The non-resonant background represents prompt production of the final state. As these two possibilities are, in principle, physically distinguishable, they must be separately gauge-invariant.

We can thus use finite propagation as a tag for identifying unstable particles without requiring that they appear in the final state. This is, after all, the way b -quarks are identified in vertex detectors. A production cross-section for an unstable particle is obtained by extracting the resonant part of the matrix element for a process containing that particle in an intermediate state and summing over all possible decay modes.

3. The process $e^+e^- \rightarrow Z^0 Z^0$

In this section we will calculate the cross-section for $e^+e^- \rightarrow Z^0 Z^0$. This is of both theoretical and practical importance. On the theoretical side it represents an example

of a calculation of the production cross-section for unstable particles. On the practical side, at high energies $e^+e^- \rightarrow Z^0 Z^0$ will be a dominant source of fermion pairs $(f_1 \bar{f}_1)$ and $(f_2 \bar{f}_2)$ due to its double resonant enhancement and hence $\sigma(e^+e^- \rightarrow Z^0 Z^0)$ is an excellent approximation to the cross-section for 4-fermion pair production. If the experimental situation warrants it, background terms can also be included without difficulty. A more detailed account can be found in ref. 3.

In the case of $e^+e^- \rightarrow f \bar{f}$, dealt with in ref.s 4,5 the invariant mass squared of the Z^0 , s , is fixed by the momenta of the incoming e^+e^- . For the process $e^+e^- \rightarrow Z^0 Z^0$ the invariant mass of the produced Z^0 's is not constant and must be somehow included in phase space integrations. It is not immediately clear how to do this and without further guidance from S -matrix theory there would seem to be considerable flexibility in how to proceed. A new ingredient is required and that is to realize that an expression for an S -matrix element can always be divided into a part that is a Lorentz-invariant function of the kinematic invariants of the problem and Lorentz-covariant objects, such as $\not{p}$ etc. The latter are known as *standard covariants*^{6,7,8,9}. It is the Lorentz invariant part that satisfies the requirements of analytic S -matrix theory and from which the resonant and non-resonant background parts are extracted while the Lorentz covariant part is untouched.

To calculate $\sigma(e^+e^- \rightarrow Z^0 Z^0)$, we begin by constructing the cross-section for $e^+e^- \rightarrow Z^0 Z^0 \rightarrow (f_1 \bar{f}_1)(f_2 \bar{f}_2)$, and will eventually sum over all fermion species. The squared invariant masses of the $f_1 \bar{f}_1$ and $f_2 \bar{f}_2$ pairs will be denoted p_1^2 and p_2^2 .

To obtain the piece of the matrix element that corresponds to finite propagation of both Z^0 's we extract the leading term in a Laurent expansion in p_1^2 and p_2^2 of the analytic Lorentz-invariant part of matrix element leaving the Lorentz-covariant part untouched. This is the doubly-resonant term and is given by

$$\begin{aligned} \mathcal{M} = & \sum_i [\bar{v}_{e^+} T_{\mu\nu}^i u_{e^-}] M_i(t, u, s_p, s_p) \\ & \times \frac{F_{ZZ}(s_p)}{p_1^2 - s_p} [\bar{u}_{f_1} \gamma^\mu (V_{Zf_L}(s_p) \gamma_L + V_{Zf_R}(s_p) \gamma_R) v_{\bar{f}_1}] \\ & \times \frac{F_{ZZ}(s_p)}{p_2^2 - s_p} [\bar{u}_{f_2} \gamma^\nu (V_{Zf_L}(s_p) \gamma_L + V_{Zf_R}(s_p) \gamma_R) v_{\bar{f}_2}] \end{aligned} \quad (3)$$

where $T_{\mu\nu}^i$ are Lorentz covariant tensors that span the tensor structure of the matrix element and γ_L, γ_R are the usual helicity projection operators. The M_i, Π_{ZZ} and V_{Zf} are Lorentz scalars that are analytic functions of the independent kinematic Lorentz invariants of the problem. F_{ZZ} is defined in the way described following eq.(1). It should be emphasized that eq.(3) is the exact form of the doubly-resonant matrix element to all orders in perturbation theory that we will now specialize to leading order.

In the procedure described above, one starts by extracting the resonant term in a scattering amplitude by Laurent expansion about the exact pole position s_p

and then specializes to lower orders by further expanding about the renormalized mass. Other authors^{10,11} have attempted to apply the techniques described above by first expanding about the renormalized mass and then adding a finite width in the denominator of the resonant part by hand. That procedure cannot be justified and leads to problems when one treats processes like $e^+e^- \rightarrow W^+W^-$. It gives rise to spurious *threshold singularities* or complex scattering angles due the production threshold's branch point being incorrectly located on the real axis.

It is important to note that Laurent expansion refers to how the resonant term is identified. The aim is only to separate $\mathcal{M}$ into gauge-invariant resonant and background pieces and there is no necessity to perform a Laurent expansion beyond its leading term although this can provide a useful way for parameterizing electroweak data¹². Although in what follows the background terms are systematically dropped, they can be kept if the experimental situation should require it.

In lowest order eq.(3) becomes, up to overall multiplicative factors,

$$\begin{aligned} \mathcal{M} = & \sum_{i=1}^2 [\bar{v}_{e^+} T_{\mu\nu}^i u_{e^-}] M_i \\ & \times \frac{1}{p_1^2 - s_p} [\bar{u}_{f_1} \gamma^\mu (V_{Zf_L} \gamma_L + V_{Zf_R} \gamma_R) v_{\bar{f}_1}] \\ & \times \frac{1}{p_2^2 - s_p} [\bar{u}_{f_2} \gamma^\nu (V_{Zf_L} \gamma_L + V_{Zf_R} \gamma_R) v_{\bar{f}_2}]. \end{aligned} \quad (4)$$

where $T_{\mu\nu}^1 = \gamma_\mu (\not{p}_{e^-} - \not{p}_1) \gamma_\nu$, $M_1 = t^{-1}$; $T_{\mu\nu}^2 = \gamma_\nu (\not{p}_{e^-} - \not{p}_2) \gamma_\mu$, $M_2 = u^{-1}$ and the final state vertex corrections take the form $V_{Zf_L} = ie\beta_L^f \gamma_L$ and $V_{Zf_R} = ie\beta_R^f \gamma_R$, The left- and right-handed couplings of the Z^0 to a fermion f are

$$\beta_L^f = \frac{t_3^f - \sin^2 \theta_W Q^f}{\sin \theta_W \cos \theta_W}, \quad \beta_R^f = -\frac{\sin \theta_W Q^f}{\cos \theta_W}.$$

Squaring the matrix element and integrating over the final state momenta for fixed p_1^2 and p_2^2 gives

$$\frac{\partial^3 \sigma}{\partial t \partial p_1^2 \partial p_2^2} = \frac{\pi \alpha^2}{s^2} (|\beta_L^e|^4 + |\beta_R^e|^4) \rho(p_1^2) \rho(p_2^2) \left\{ \frac{t}{u} + \frac{u}{t} + \frac{2(p_1^2 + p_2^2)^2}{ut} - p_1^2 p_2^2 \left(\frac{1}{t^2} + \frac{1}{u^2} \right) \right\} \quad (5)$$

with

$$\begin{aligned} \rho(p^2) &= \frac{\alpha}{6\pi} \sum_f (|\beta_L^f|^2 + |\beta_R^f|^2) \frac{p^2}{|p^2 - s_p|^2} \theta(p_0) \theta(p^2) \\ &\approx \frac{1}{\pi} \cdot \frac{p^2 (\Gamma_Z/M_Z)}{(p^2 - M_Z^2)^2 + \Gamma_Z^2 M_Z^2} \theta(p_0) \theta(p^2) \end{aligned}$$

where the sum is over fermion species. Note that $\rho(p^2) \rightarrow \delta(p^2 - M_Z^2) \theta(p_0)$ as $\text{Im}(s_p) \rightarrow 0$ which is the result obtained by cutting a free propagator. The variables s , t , u ,

p_1^2 and p_2^2 in eq.(5) arise from products of standard covariants and external wave functions and therefore take real values dictated by the kinematics.

Integrating over t , p_1^2 and p_2^2 leads to

$$\sigma(s) = \int_0^s dp_1^2 \int_0^{(\sqrt{s}-\sqrt{p_1^2})^2} dp_2^2 \sigma(s; p_1^2, p_2^2) \rho(p_1^2) \rho(p_2^2), \quad (6)$$

where

$$\begin{aligned} \sigma(s; p_1^2, p_2^2) &= \frac{2\pi\alpha^2}{s^2} (|\beta_L^e|^4 + |\beta_R^e|^4) \\ &\times \left\{ \left(\frac{1 + (p_1^2 + p_2^2)^2/s^2}{1 - (p_1^2 + p_2^2)/s} \right) \ln \left(\frac{-s + p_1^2 + p_2^2 + \lambda}{-s + p_1^2 + p_2^2 - \lambda} \right) - \frac{\lambda}{s} \right\} \end{aligned}$$

and $\lambda = \sqrt{s^2 + p_1^4 + p_2^4 - 2sp_1^2 - 2sp_2^2 - 2p_1^2p_2^2}$. For $p_1^2 = p_2^2 = M_Z^2$ this agrees with known results¹³.

4. The Process $e^+e^- \rightarrow \bar{\nu}_e e^- u \bar{d}$

Massless particles, such as the photon, are always problematic for analytic S -matrix analyses. In this section the process $e^+e^- \rightarrow \bar{\nu}_e e^- u \bar{d}$ will be considered with particular emphasis on how take into account the finite width of the W boson while at the same time maintaining QED gauge invariance. The dominant contribution to this process is expected to come from physical W^+ production in which $e^+e^- \rightarrow \bar{\nu}_e e^- W^+ \rightarrow \bar{\nu}_e e^- u \bar{d}$ via diagrams of the type shown in Fig.1. Assume for the moment that the W is stable and therefore has the usual propagator $\sim (p^2 + M_W^2)^{-1}$. It is expected that the matrix element for this process should satisfy the usual condition for QED gauge-invariance

$$q_\mu G^\mu = 0 \quad (7)$$

where G^μ is the matrix element with the external electromagnetic current and photon propagator amputated and q_μ is the momentum in the photon propagator. When this test is applied to the diagrams of Fig.1. one finds that (7) does not hold. The reason is that there exist other Feynman diagrams that produce the same final state and only when these are included is the condition (7) correctly satisfied¹⁴. However the additional diagrams that must be included do not contain internal W 's that could go on-mass-shell and are therefore expected to be subdominant.

It might be thought that gauge dependence in subdominant terms of no concern but for the process in question it plays a crucial rôle. It can be shown¹⁴ that as $q^2 \rightarrow 0$ the resulting cross-section behaves as $\sim 1/q^2$ provided (7) is satisfied and $\sim 1/q^4$ when it is not. Without the maintenance of QED gauge invariance the cross-section then blows up too strongly as $q^2 \rightarrow 0$.

The inclusion of subdominant diagrams to cancel the gauge-dependence in the dominant contribution contrasts sharply with what was done in section 3. The Laurent expansion extracts an exactly gauge-invariant resonant term and eliminates from




Figure 1: Tree-level diagrams contributing to the resonant part of the process $e^+e^- \rightarrow \bar{\nu}_e e^- W^+ \rightarrow \bar{\nu}_e e^- u \bar{d}$. (courtesy of the authors of ref.14)

it any non-resonant (subdominant) pieces. These pieces that are removed may be ultimately combined with other contributions to yield an overall gauge-invariant non-resonant contribution if required.

Now consider the case when the finite width of the W boson in the diagrams of Fig.1 is to be taken into account. If the procedure described in sections 2 and 3 is applied to the process $e^+e^- \rightarrow \bar{\nu}_e e^- W^+ \rightarrow \bar{\nu}_e e^- u \bar{d}$ to extract the dominant resonant piece then the resulting matrix element does not satisfy the condition (7) and the resulting cross-section blows up too strongly as $q^2 \rightarrow 0$. A number of solutions have been proposed and their relative merits examined in ref. 14. One that has received much attention is that of Baur and Zeppenfeld¹⁵. Their solution is to add the imaginary parts of certain higher-order diagrams, specifically the 1-loop fermionic correction to the $W^+W^-\gamma$ vertex, which they show exactly removes the QED gauge dependence. This method has the practical disadvantage that it requires an additional calculation that is significantly more complicated to obtain than the rest of the tree-level amplitude. Moreover once it is calculated it is immediately largely canceled by the gauge-dependent terms in the resonant part.

In this section we describe an alternative more economical approach that lies more in line with the strategies adopted in section 3. Instead of seeking additional higher-order gauge-dependent corrections to cancel the QED gauge dependence, we eliminate it by means of a projection operator under which the exact matrix element is known to be invariant.

Suppose the approach of section 3 has been used to obtain the leading resonant contribution to the process $e^+e^- \rightarrow \bar{\nu}_e e^- W^+ \rightarrow \bar{\nu}_e e^- u \bar{d}$. After amputating the external electromagnetic current and photon propagator a Green's function, G_μ , is obtained that can be written in terms of a set of standard covariants, l_i^μ , and associated Lorentz scalar functions, $A_i(p_1^2, p_2^2, \dots)$,

$$G^\mu = \sum_i l_i^\mu(p_1, p_2, \dots) A_i(p_1^2, p_2^2, \dots) \quad (8)$$

where the arguments p_1, p_2 , etc. are 4-momenta that appear in the problem. The l_i^μ , of course, must transform in the same way that G^μ does.

The gauge condition on Green's functions $q_\mu G^\mu = 0$ means that the A_i are not linearly independent since they satisfy the condition $\sum_i (q \cdot l_i) A_i = 0$.

Gauge-invariant Green's functions can be obtained using the methods of Bardeen and Tung¹⁶. Suppose that we have some exact Green's function expressed in terms of standard covariants and Lorentz scalars as in eq.(8). Since $q_\mu G^\mu = 0$ the Green's function is invariant under the action of the operator

$$I_\nu^\mu = g_\nu^\mu - \frac{p^\mu q_\nu}{p \cdot q} \quad (9)$$

where p is some conveniently chosen momentum. The spinor-Lorentz tensor structure of the Green's function is spanned by the set $\{I_\nu^\mu l_i^\nu\}$. That is

$$G^\mu(p_1, p_2, \dots) = \sum_i (I_\nu^\mu l_i^\nu(p_1, p_2, \dots)) A_i(p_1^2, p_2^2, \dots). \quad (10)$$

For an approximate Green's function that does not satisfy the QED gauge-condition (7), the operator I_ν^μ serves as a projection operator onto $\{I_\nu^\mu l_i^\nu\}$. From the $\{I_\nu^\mu l_i^\nu\}$ it is always possible to construct a new basis that is free of kinematic singularities that might have occurred when, for example, $(p \cdot q) = 0$. In practice, however, it is simpler to just choose the vector p such that $(p \cdot q)$ is not small in the kinematic region of interest.

In general consider a Green's function G^μ calculated using some incomplete expansion up to some order. It will consist of two parts

$$G^\mu = G_0^\mu + G_1^\mu \quad (11)$$

where G_0^μ is a consistent gauge-invariant contribution correct to given order of the calculation. G_1^μ is a spurious higher-order gauge-dependent correction. Because G_0^μ satisfies the gauge condition, $q_\mu G_0^\mu = 0$, it is invariant under the action of the operator I_μ^ν . Thus at a given order we may make the replacement

$$G^\mu \rightarrow I_\nu^\mu G^\nu = G_0^\mu + I_\nu^\mu G_1^\nu.$$

Only G_1^μ is affected by the projection operator but since it is of higher order this is of no concern. The important point is that it now satisfies the gauge condition $q_\mu (I_\nu^\mu G_1^\nu) = 0$ and is prevented from causing too strong a numerical blow up as $q^2 \rightarrow 0$.

Now, of course, the result depends on the choice of the vector, p , which when judiciously done can be used to significantly simplify calculations. The dependence on p , as it appears only in higher-orders is analogous to renormalization scheme dependence, or the μ -dependence in the $\overline{\text{MS}}$ scheme, that is an inescapable feature of any perturbative expansion. As with scheme dependence, we can be sure that it will become ever weaker as higher orders included since the exact Green's function is invariant under the action of I_ν^μ .

Details of applying the projection operator to the process $e^+e^- \rightarrow \bar{\nu}_e e^- W^+ \rightarrow \bar{\nu}_e e^- u \bar{d}$ may be found in ref. 17 but may be summarized as follows:

The methods of section 3 are used to obtain the resonant part of the matrix element $\mathcal{M}$ which takes the form

$$\mathcal{M} = J_\lambda^\gamma P_\mu^\lambda G^\mu \quad (12)$$

where J_λ^γ is the electromagnetic current of the electron and P_μ^λ is the photon propagator. It is simply a matter of inserting the projection operator, (9), to obtain

$$\mathcal{M} = J_\lambda^\gamma P_\nu^\lambda (I_\mu^\nu G^\mu). \quad (13)$$

The terms in parentheses satisfy the gauge-condition (7) and therefore the cross-section formed from $\mathcal{M}$ behaves $\sim 1/q^2$ as $q^2 \rightarrow 0$. Terms eliminated by the projection operator must cancel in the exact matrix element.

1. R. E. Peierls, *Proceedings of the 1954 Glasgow Conference on Nuclear and Meson Physics*, Pergamon Press, New York, (1955) 296.
2. M. Veltman, *Physica* **29** (1963) 186.
3. R. G. Stuart, hep-ph/9504215.
4. R. G. Stuart, *Phys. Lett. B* **262** (1991) 113.
5. R. G. Stuart, *Phys. Rev. Lett.* **70** (1993) 3193.
6. A. C. Hearn, *Nuovo Cimento* **21** (1961) 333.
7. K. Hepp, *Helv. Phys. Acta* **36** (1963) 355.
8. D. N. Williams, preprint UCRL-11113 (1963).
9. K. Hepp, *Helv. Phys. Acta* **37** (1964) 11.
10. A. Aeppli, F. Cuyper and G. J. van Oldenborgh, *Phys. Lett. B* **314** (1993) 413;
11. A. Aeppli, G. J. van Oldenborgh and D. Wyler, *Nucl. Phys. B* **428** (1994) 126.
12. R. G. Stuart, hep-ph/9602300.
13. R. W. Brown and K. O. Mikaelian, *Phys. Rev. D* **19** (1979) 922.
14. E. N. Argyle *et al.*, *Phys. Lett. B* **358** (1995) 339.
15. U. Baur and D. Zeppenfeld, *Phys. Rev. Lett.* **75** (1995) 1002.
16. W. A. Bardeen and W.-K. Tung, *Phys. Rev.* **173** (1968) 1423.
17. R. G. Stuart, hep-ph/9603351.